arXiv:1205.2521v1 [math.PR] 11 May 2012

From Minority Game to Black&Scholes pricing

Matteo Ortisi[*], Valerio Zuccolo[†]

Abstract

In this paper we study the continuum time dynamics of a stock in a market where agents behavior is modeled by a Minority Game with number of strategies for each agent $S = 2$ and "fake" market histories. The dynamics derived is a generalized geometric Brownian motion; from the Black&Scholes formula the calibration of the Minority Game, by means of the game parameter σ^2, on the European options on DAX Index market is performed. An "(α, σ^2) -matrix" containing, given options' moneyness and maturities, values of the parameters α and σ^2 that make the theoretical option price agree with the market price is constructed. We conclude that the asymmetric phase of the Minority Game with α close to α_c is coherent with options implied volatility market.

Key words: Minority Game, Agent-based models, Option pricing, Market calibration.

1 Introduction

In recent years researchers endeavored to build models that reproduce some empirical statistical regularities of the real financial markets, such as volatility clusters, fat tails, scaling, occurrence of crashes, etc., known as "stylized facts" (see [10, 12, 14]). One possible way to address this problem is the "black-box" stochastic approach. A complex stochastic process which possesses the relevant characteristics of the desired empirical facts is constructed. Process used in this kind of models are the nonlinear diffusion models, the Lévy processes, the jump processes and stochastic volatility models. Another possibility is to follow the agent-based framework, which models the market and derives from the interaction between players the stock price dynamics. Herd Behavior and Minority Game are examples of these models. Choosing between these two models families relies on the purpose of the modeling itself. "Black-box" stochastic modeling is used in financial mathematics, where the main goal is the practical use, i.e. pricing of derivatives or portfolio allocation. Agent-based models are mainly used in economics where the theoretical aspects, the explanation and understanding of financial markets, are the main intent.

There is a complementarity between these two approaches; indeed "black-box" stochastic models lack of explanatory power by construction and agent-based models can be hardly used for pricing purposes given their complexity. Moreover one of most attractive features of the usual Black&Scholes type models is the possibility of obtaining closed, exact formulas for the pricing of financial derivative securities. Such qualities are fundamental from the point of view of finance practitioners, like financial institutions, that require fast calculations, closed pricing formula for calibration purposes and tools to hedge risky assets.

---

[*] Pioneer Investments, Galleria San Carlo 6, 20122 Milano, Italy (matteo.ortisi@gmail.com)

[†] Intesa Sanpaolo, Piazza Paolo Ferrari 10, 20100 Milano, Italy (valerio.zuccolo@polimi.it)

A very promising way to combine these main issues, is to focus on highly simplified toy models of financial markets relying on the Minority Game (MG) [4]. Variants of this model have been shown to reproduce quite accurately such stylized facts of financial markets (see [1, 3, 5, 11, 20]) and moreover the continuum time limit of the MG provides a bridge between the adherence to empirically observed features of real markets and the practical usability of "black-box" stochastic models, since its evolution follows a system of stochastic differential equations (see [15], [8]).

In this framework, the stock price process is driven by the excess demand or overall market bid. The excess demand evolution process for real and "fake" market histories has been explicitly obtained in [8]. While a game with real market histories can fit stock prices better than a game with "fake" ones, the latter is more mathematically tractable, especially if the purpose is to obtain closed formulas and not run the game through simulations. Models with "fake" market histories, where at each point in time all agents are given random rather than real market data upon which to base their decisions, have the advantage of being Markovian and hence suitable to the application of classic results, like Girsanov theorem, that are at the hearth of option pricing techniques in mathematical finance. On the other side, models with real market histories are strongly non-Markovian and a dynamics simple like (3) is no more available. The dependence of the process on the history makes meaningless all the averages present in equation (3); the evolution process becomes a system of stochastic differential equations whose parameters are to be obtained as a solution of a dynamic system defining the bid evolution process (see [8] for details). Since our aim is to calibrate the MG on the real market of European call options, as a first attempt to obtain a closed formula for pricing an European call option on a single stock it is easier, and maybe compulsory, to use the simplest version of the minority game with "fake" market histories and to recover the excess demand directly from the scores difference stochastic differential equations obtained in [15]. Using this approach it is possible to apply the usual "change of measure technique", that guarantee the absence of arbitrage in the model and the existence of the hedging portfolio, and hence to obtain closed pricing formulas allowing the model calibration to the market. Undoubtedly the MG with real market history must be investigated and the approach used in [8] must be kept in mind for further development.

The work is organized as follow. In Section 2 the classical MG model and its continuum time limit version are described. In Section 3 we derive the stock price dynamics and in Section 4, by using the classical risk-neutral pricing techniques (see [19]), we apply the Black&Scholes formula to price European options. Section 5 contains the calibration of the MG on the European options market; finding out values of α and σ that make the theoretical price of an option on DAX Index obtained via Black&Scholes formula agrees with the market price, an "(α, σ^2)-matrix" with options' moneynees and maturity as entries is constructed. In Section 6 we report our conclusions.

2 The Model

In this section the basic concept and the main results of MG useful for our purposes will be exposed (for a comprehensive introduction to MG refer to [6, 9]). Here we are considering a slightly modified version of the MG, where there are N agents and its dynamics is defined in terms of dynamical variables $U_{s,i}(t)$, scores corresponding to each of the possible agents strategy choices $s = +w, -w$.

Each agent takes a decision (strategy choice) $s_i(t)$ with

$$Prob\{s_i(t) = s\} = \frac{e^{\Gamma_i U_{s,i}(t)}}{\sum_{s'} e^{\Gamma_i U_{s',i}(t)}}$$

where $\Gamma_i > 0$, and $s' \in \{-w, +w\}$. The original MG corresponds to $\Gamma_i = \infty$ and was generalized to $\Gamma_i = \Gamma < \infty$ [2]; here we consider the latter case.

The public information variable $\mu(t)$, that represent the common knowledge of the past record, is given to all agents; it belongs to the set of integers $\{1, \ldots, P\}$ and can either be the binary encoding of last M winning choices or drawn at random from a uniform distribution:

$$Prob\{\mu(t) = \mu\} = \frac{1}{P}, \quad \mu = 1, \ldots, P.$$

Here we consider the latter case (see [2]).

The strategies $a^\mu_{s,i}$ are uniform random variables taking values $\pm w$ ($Prob\{a^\mu_{s,i} = \pm w\} = 1/2$) independent on i, s and μ. Here we consider $S = 2$, i.e. 2 strategy for each agent that are randomly drawn at the beginning of the game and kept fixed.

On the basis of the outcome $B(t) = \sum_{i=1}^N a^{\mu(t)}_{s_i(t),i}$ each agent update his scores according to

$$U_{s,i}(t + \delta t) = U_{s,i}(t) - a^{\mu(t)}_{s,i} \frac{B(t)}{P}, \tag{1}$$

where $\delta t \to 0$ is the time increment.

Let us introduce the following random variables (to ease the notation the choices $+w$ and $-w$ are shorted with $+$ and $-$)

$$\xi^\mu_i = \frac{a^\mu_{+,i} - a^\mu_{-,i}}{2}, \quad \Theta^\mu = \sum_{i=1}^N \frac{a^\mu_{+,i} + a^\mu_{-,i}}{2}$$

and their averages

$$\overline{\xi_i \Theta} = \frac{1}{P} \sum_{\mu=1}^P \xi^\mu_i \Theta^\mu, \quad \overline{\xi_i \xi_j} = \frac{1}{P} \sum_{\mu=1}^P \xi^\mu_i \xi^\mu_j.$$

The only relevant quantity in the dynamics is the difference between the scores of the two strategies:

$$y_i(t) = \Gamma \frac{U_{+,i}(\tau) - U_{-,i}(\tau)}{2}, \tag{2}$$

where $\tau = \frac{t}{\Gamma}$.

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be the probability space respect to which all our random variables are defined, $y = (y_i)_{1 \le i \le N}$, $\Theta = (\Theta^\mu)_{1 \le \mu \le P}$ and $\xi = (\xi_i)_{1 \le i \le N}$.

As shown in [15], if $P/N = \alpha \in \mathbb{R}_+$, $S = 2$ and $\Gamma_i = \Gamma > 0$ for all i, the dynamics of the continuum time limit of the MG is given by the following N-dimensional stochastic differential equation

$$dy_i(t) = \left(-\overline{\xi_i \Theta} - \sum_{j=1}^N \overline{\xi_i \xi_j} \tanh(y_j(t)) \right) dt + A_i(y, \alpha, N, \Gamma, \xi) d\mathbf{W}(t), \quad i = 1, \ldots, N \tag{3}$$

where

$\mathbf{W}(t)$ is an N-dimensional Wiener process,

A_i is the i-th row of the $N \times N$ matrix $A = (A_{ij})$ such that

$$(AA')_{ij}(y, \alpha, N, \Gamma, \xi) = \frac{\Gamma \sigma_N^2(y)}{\alpha N} \overline{\xi_i \xi_j}.$$

The dynamics of σ_N^2 is well known; the relevant feature for our purposes is its limit behavior, as N grows to infinity, with respect to α: for $\alpha \geq \alpha_c$, $\lim_{N \to \infty} \frac{\sigma_N^2(y)}{N} \leq 1$, $\forall y$, while for $\alpha < \alpha_c$, $\lim_{N \to \infty} \frac{\sigma_N^2(y)}{N^2} \leq k$, $\forall y$ and with k constant. For more details see [9].

With respect to the usual version of the MG where $s = +1, -1$, here each agent gives each buy or sell decision the same weight w. The introduction of parameter w, that, as we will see, improves the calibration of the game on the options market, affect the scores updating (1) of a factor w^2, coming from $B(t)$ and $a^{\mu}_{s,i}$, and hence the variable y defined in (2) of the same factor. This reflects on dynamics (3) where both the drift term and the AA' elements are multiplied (with respect to the case $s = \pm 1$) by the same w^2 factor. This does not affect the fluctuations (i.e. σ_N^2) and the relation between σ_N^2 and α, but, as we will see in the sequel, the stock price dynamics and the expression of the option implied volatility, fundamental in the calibration process.

3 Stock price dynamics

In this section, using the scores difference stochastic differential equations, the overall market bid and hence the stock price dynamics is derived.

Let us consider a single stock in a market modeled by the continuum time MG; following [5], we define the stock price dynamics in terms of excess demand, as

$$\log p(t + \delta t) = \log p(t) + \frac{B(t)}{N} \frac{1}{w} \tag{4}$$

(price dynamics does not depend on the weight w of the bet).

Since dynamics (4) has a deep impact on the results we will obtain, it worths to spend few words to explain our choice.

Defining the stock price dynamics as in (4), we are assuming the volume N to be constant. Even if a choice of fluctuating volumes $N(t)$ would be more realistic, it would imply a new derivation of the time-continuous dynamics (3) that would become a system of a time dependent number of stochastic differential equations with terms like $\overline{\xi_i \Theta}$ now time dependent. This would have a heavy impact: all the results obtained in the limit as N grows to infinity should be re-obtained in terms of uniform convergence over t. If they still hold, also proposition (4.1), crucial for our purposes since proves the existence of a risk neutral measure, must be re-derived; this could be even harder since it relies on the Girsanov theorem: does exist a version of such theorem for a $N(t)$ dimensional stochastic process? Since the present one is, as far as we know, the first attempt to obtain a closed option pricing formula starting from a MG, we think that it is better to stick to the simplest case, and then to move toward more realistic, and mathematically complicated, frameworks.

Our aim is to obtain the continuum time dynamics of the stock price $p(t)$ from the continuum time dynamics of $\mathbf{y}(t)$ defined by (2). The key point is the following identity, that can be obtained directly from (1)

$$\frac{U_{+,i}(t + \delta t) - U_{-,i}(t + \delta t)}{2} = \frac{U_{+,i}(t) - U_{-,i}(t)}{2} - \frac{a^{\mu(t)}_{+,i} - a^{\mu(t)}_{-,i}}{2} \frac{B(t)}{P}.$$

Taking into account relation (2) and that Γ is finite, we obtain

$$\frac{y_i(\Gamma t)}{\Gamma} - \frac{y_i(\Gamma(t+\delta t))}{\Gamma} = -\xi_i^{\mu(t)} \frac{B(t)}{\alpha N}.$$

In the continuum time limit $\delta t \to 0$,

$$dy_i(\Gamma t) = -\Gamma \xi_i^{\mu(t)} \frac{B(t)}{\alpha N}, \quad \forall i = 1, 2, \ldots, N. \tag{5}$$

Dynamics (5) reflects the stock price dynamics (4), where $\frac{B(t)}{N}$ is the generator of $\log p(t)$; indeed, like the right hand side of (4), also the right hand side of (5) does not contain a dt term and $-\Gamma \xi_i^{\mu(t)} \frac{B(t)}{\alpha N}$ is the generator of $y_i(\Gamma t)$.

Multiplying both sides of (5) by $\xi_i^{\mu(t)}$, averaging over all the agents i, and taking the limit as N goes to infinity we have

$$\lim_{N\to\infty} \frac{1}{N} \sum_{i=1}^{N} \xi_i^{\mu(t)} dy_i(\Gamma t) = -\frac{\Gamma}{\alpha} \lim_{N\to\infty} \frac{1}{N} \sum_{i=1}^{N} \left(\xi_i^{\mu(t)}\right)^2 \frac{B(t)}{N}.$$

Since the $\xi_i^{\mu(t)}$ are independent from $B(t)$ (the difference between two actions given the past history $\mu(t)$ is fixed at the beginning of the game and does not depend on the action chosen by the agent on the basis of the score function $U_{s,i}(t)$),

$$\lim_{N\to\infty} \frac{1}{N} \sum_{i=1}^{N} \left(\xi_i^{\mu(t)}\right)^2 \frac{B(t)}{N} = \mathbb{E}\left[\left(\xi_i^{\mu(t)}\right)^2\right] \lim_{N\to\infty} \frac{B(t)}{N} = \frac{w^2}{2} w d \log p(t);$$

it follows that

$$d \log p(t) = -\frac{2\alpha}{w^3 \Gamma} \lim_{N\to\infty} \frac{1}{N} \sum_{i=1}^{N} \xi_i^{\mu(t)} dy_i(\Gamma t). \tag{6}$$

Integrating over $[0, t]$ both sides of (6) and taking into account equation (3), it follows that

$$\begin{aligned} p(t) \quad &= \quad p(0) \exp\left\{ \lim_{N\to\infty} \int_0^t \frac{2\alpha}{w^3 N} \sum_{i=1}^{N} \left[\xi_i^{\mu(s)} \left(\overline{\xi_i \Theta} + \sum_{j=1}^{N} \overline{\xi_i \xi_j} \tanh(y_j(\Gamma s)) \right) \right] ds \right. \\ &\quad - \left. \lim_{N\to\infty} \int_0^t \frac{2\alpha}{w^3 N \Gamma} \sum_{i=1}^{N} \left(\xi_i^{\mu(s)} A_i \right) d\mathbf{W}(\Gamma s) \right\}. \end{aligned} \tag{7}$$

The following proposition on the drift and diffusion term of equation (7) holds:

Proposition 3.1 *Let $t = O(N)$, that is $\lim_{N\to+\infty} \frac{t}{N} = k$. The drift term of (7) is at most $O(N)$; the diffusion term is different from zero $\forall \alpha$, finite for $\alpha \geq \alpha_c$ and at most $O(N)$ for $\alpha < \alpha_c$.*

Proof

It is easy to see that, by applying the Law of Large Numbers, $\forall i \lim_{N\to\infty} \overline{\xi_i^2} = \frac{w^2}{2}$, $\lim_{N\to\infty} \overline{\xi_i \xi_j} = 0$ and $\lim_{N\to\infty} \overline{\xi_i \Theta} = 0$ (for detailed computations see [18]); it follows that

$$\lim_{N\to\infty} \frac{2\alpha}{w^3 N} \sum_{i=1}^{N} \left[\xi_i^{\mu(s)} \left(\overline{\xi_i \Theta} + \sum_{j=1}^{N} \overline{\xi_i \xi_j} \tanh(y_j(\Gamma s)) \right) \right] = \frac{\alpha}{w^3} \lim_{N\to\infty} \frac{1}{N} \sum_{i=1}^{N} \xi_i^{\mu(s)} \tanh(y_i(\Gamma s)) < k\alpha,$$

with k finite constant. Hence the drift term is at most $k\alpha N$ and, due to different initial conditions and the stochastic evolution of y_i, in general different from zero.

Consider the zero mean random variable

$$Y = \int_0^t \frac{2\alpha}{w^3 N\Gamma} \sum_{i=1}^{N} \left(\xi_i^{\mu(s)} A_i\right) d\mathbf{W}(\Gamma s);$$

since

$$\frac{2\alpha}{w^3 N\Gamma} \sum_{i=1}^{N} \left(\xi_i^{\mu(s)} A_i\right) d\mathbf{W}(\Gamma s) = \sum_{j=1}^{N} \left(\frac{2\alpha}{w^3 N\Gamma} \sum_{i=1}^{N} \xi_i^{\mu(s)} A_{ij} \right) dW_j(\Gamma s),$$

Y has variance

$$\begin{aligned} \nu &= \int_0^t \sum_{j=1}^{N} \left(\frac{2\alpha}{w^3 N\Gamma} \sum_{i=1}^{N} \xi_i^{\mu(s)} A_{ij} \right)^2 d(\Gamma s) \\ &= \int_0^t \frac{4\alpha^2}{w^6 N^2 \Gamma} \sum_{i=1}^{N} \left(\xi_i^{\mu(s)}\right)^2 (AA')_{ii} ds + \int_0^t \frac{4\alpha^2}{w^6 N^2 \Gamma} \sum_{i,j=1, i\neq j}^{N} \xi_i^{\mu(s)} \xi_j^{\mu(s)} (AA')_{ij} ds. \end{aligned} \quad (8)$$

As N grows to infinity, by applying the Law of Large Numbers, $\lim_{N\to\infty} AA' = \lim_{N\to\infty} \frac{\Gamma\sigma_N^2}{2\alpha N} w^2 I$, where I is the identity $N \times N$ matrix (see [18] for details), and the first term of the right hand side of (8) is equal to $\lim_{N\to\infty} \frac{\alpha\sigma_N^2 t}{w^2 N^2}$, while the second one is equal to 0. Since for $\alpha \geq \alpha_c$, $0 < \frac{\sigma_N^2}{N} < 1$ and for $\alpha < \alpha_c$, $0 < \frac{\sigma_N^2}{N} < kN$, the thesis follows.

□

It is worth to note that the hypothesis $t = O(N)$ is necessary to have the diffusion term different from zero and that it is the proportionality factor of the time needed to reach the stationary state of the MG. In numerical terms it means that, given a maturity t, a higher number of player needs a higher number of time steps, i.e. the player must interact more times to reach an equilibrium.

Consider the dynamics

$$\begin{aligned} q(t) &= p(0) \exp \left\{ \int_0^t \frac{2\alpha}{w^3 N} \sum_{i=1}^{N} \left[\xi_i^{\mu(s)} \left(\overline{\xi_i \Theta} + \sum_{j=1}^{N} \overline{\xi_i \xi_j} \tanh(y_j(\Gamma s)) \right) \right] ds \right. \\ &\quad - \left. \int_0^t \frac{2\alpha}{w^3 N\Gamma} \sum_{i=1}^{N} \left(\xi_i^{\mu(s)} A_i\right) d\mathbf{W}(\Gamma s) \right\}; \end{aligned}$$

by definition $q(t)$ converges pointwise towards $p(t)$ ($\lim_{N\to\infty} q(t) = p(t)$, $\forall \omega \in \Omega$) and hence almost surely; it follows that a.s. $\forall \epsilon > 0$ there exists $\overline{N} > 0$ such that, $\forall N > \overline{N}$, $|p(t) - q(t)| < \epsilon$. By assuming $t = kN$ and N finite but sufficiently large, we have that a.s. $q(t) \sim p(t)$ and proposition (3.1) holds for $q(t)$.

From now on we assume $t = kN$ and N finite but sufficiently large to have

$$\begin{aligned} p(t) &= p(0) \exp \left\{ \int_0^t \frac{2\alpha}{w^3 N} \sum_{i=1}^{N} \left[\xi_i^{\mu(s)} \left(\overline{\xi_i \Theta} + \sum_{j=1}^{N} \overline{\xi_i \xi_j} \tanh(y_j(\Gamma s)) \right) \right] ds \right. \\ &\quad - \left. \int_0^t \frac{2\alpha}{w^3 N\Gamma} \sum_{i=1}^{N} \left(\xi_i^{\mu(s)} A_i\right) d\mathbf{W}(\Gamma s) \right\}. \end{aligned} \quad (9)$$

Let us define $\mathcal{G}_t = \sigma(W(\Gamma s), \xi^{\mu(s)}; s \leq t)$ the filtration generated by both the processes $\mathbf{W}$ and ξ. Since the drift and the diffusion terms are just the exponential of the drift and the diffusion of equation (3) multiplied by $\xi_i^{\mu(s)}$, they are adapted to the filtration $\mathcal{G}_t$.

The differential of the stock price process is therefore, by Ito's formula,

$$\begin{aligned}
\frac{dp(t)}{p(t)} &= \left[\frac{2\alpha}{w^3 N}\sum_{i=1}^{N}\xi_i^{\mu(t)}\left(\overline{\xi_i\Theta}+\sum_{j=1}^{N}\overline{\xi_i\xi_j}\tanh(y_j(\Gamma t))\right)\right. \\
&\quad \left.+\frac{2\alpha^2}{\Gamma w^6 N^2}\left(\sum_{i=1}^{N}\xi_i^{\mu(t)}A_i\right)\left(\sum_{i=1}^{N}\xi_i^{\mu(t)}A_i\right)'\right]dt-\frac{2\alpha}{w^3 N\Gamma}\sum_{i=1}^{N}\left(\xi_i^{\mu(t)}A_i\right)d\mathbf{W}(\Gamma t) \\
&= \left[\frac{2\alpha}{w^3 N}\sum_{i=1}^{N}\xi_i^{\mu(t)}\left(\overline{\xi_i\Theta}+\sum_{j=1}^{N}\overline{\xi_i\xi_j}\tanh(y_j(\Gamma t))\right)+\frac{2\alpha^2}{w^6\Gamma N^2}\sum_{j=1}^{N}\left(\sum_{i=1}^{N}\xi_i^{\mu(t)}A_{ij}\right)^2\right]dt \\
&\quad -\frac{2\alpha}{w^3 N\Gamma}\sum_{i=1}^{N}\left(\xi_i^{\mu(t)}A_i\right)d\mathbf{W}(\Gamma t).
\end{aligned}$$

4 Derivative security pricing

In this section we develop the risk-neutral pricing of a derivative security on a single stock whose price dynamics is given by (9). We follow the usual scheme: construction of the risk-neutral measure, relying on Girsanov Theorem, for the discounted process and, by using the Martingale Representation Theorem, of the replication portfolio that allows to hedge a short position in the derivative security (see [19], p. 209-220). At the end of the section we apply Black&Scholes formula to price a European call option.

4.1 Discounted stock dynamics under risk-neutral

Consider the interest rate process $R(s)$ adapted to the filtration $\mathcal{G}_t$, $0 \leq t \leq T$. The discount process $D(t) = e^{-\int_0^t R(s)ds}$ has, by Ito's formula, differential $dD(t) = -R(t)D(t)dt$.

The discounted stock process is

$$\begin{aligned}
D(t)p(t) &= p(0)\exp\left\{\int_0^t -R(s)+\frac{2\alpha}{w^3 N}\sum_{i=1}^{N}\left[\xi_i^{\mu(s)}\left(\overline{\xi_i\Theta}+\sum_{j=1}^{N}\overline{\xi_i\xi_j}\tanh(y_j(\Gamma s))\right)\right]ds\right. \\
&\quad \left.-\int_0^t\frac{2\alpha}{w^3 N\Gamma}\sum_{i=1}^{N}\left(\xi_i^{\mu(s)}A_i\right)d\mathbf{W}(\Gamma s)\right\},
\end{aligned}$$

and its differential

$$\begin{aligned}
\frac{d(D(t)p(t))}{D(t)p(t)} &= \left[\frac{2\alpha}{w^3 N}\sum_{i=1}^{N}\left[\xi_i^{\mu(s)}\left(\overline{\xi_i\Theta}+\sum_{j=1}^{N}\overline{\xi_i\xi_j}\tanh(y_j(\Gamma s))\right)\right]\right. \\
&\quad \left.+\frac{2\alpha^2}{w^6\Gamma N^2}\sum_{j=1}^{N}\left(\sum_{i=1}^{N}\xi_i^{\mu(t)}A_{ij}\right)^2-R(t)\right]dt-\frac{2\alpha}{w^3 N\Gamma}\sum_{i=1}^{N}\left(\xi_i^{\mu(t)}A_i\right)d\mathbf{W}(\Gamma t).
\end{aligned}$$

Proposition 4.1 *If $T = O(N)$ there exists a probability measure $\widetilde{\mathbb{P}}$ (risk neutral measure) under which the discounted stock price $D(t)p(t)$ is a $\widetilde{\mathbb{P}}$-martingale.*

Proof

Let us define the market price of risk equation to be

$$\begin{aligned}\frac{2\alpha}{w^3 N\Gamma}\sum_{i=1}^{N}\left(\xi_i^{\mu(s)}A_i\right)\gamma(\Gamma t) &= \left[\frac{2\alpha}{w^3 N}\sum_{i=1}^{N}\left[\xi_i^{\mu(s)}\left(\overline{\xi_i\Theta}+\sum_{j=1}^{N}\overline{\xi_i\xi_j}\tanh(y_j(\Gamma s))\right)\right]+\right.\\ &\left.+\frac{2\alpha^2}{w^6\Gamma N^2}\sum_{j=1}^{N}\left(\sum_{i=1}^{N}\xi_i^{\mu(t)}A_{ij}\right)^2-R(t)\right],\end{aligned}$$

where $\gamma(\Gamma t) = (\gamma_1(\Gamma t),\ldots,\gamma_N(\Gamma t))$ is an unknown process; the differential of the discounted process becomes

$$d(D(t)p(t)) = D(t)p(t)\frac{2\alpha}{w^3 N\Gamma}\sum_{i=1}^{N}\left(\xi_i^{\mu(s)}A_i\right)[\gamma(\Gamma t)dt - d\mathbf{W}(\Gamma t)].$$

Let $||\cdot||$ denote the usual Euclidean norm; we show that for $T = O(N)$, $\int_0^T ||\gamma(\Gamma u)||^2 d(\Gamma u) < \infty$. Consider N finite and sufficiently large (we remind that proposition (3.1) for N finite and sufficiently large); since, by proposition (3.1)

$$\int_0^T\left|\left|\frac{2\alpha}{w^3 N\Gamma}\sum_{i=1}^{N}\left(\xi_i^{\mu(u)}A_i\right)\right|\right|^2 d(\Gamma u) = \int_0^T\sum_{j=1}^{N}\left(\frac{2\alpha}{w^3 N\Gamma}\sum_{i=1}^{N}\xi_i^{\mu(u)}A_{ij}\right)^2 d(\Gamma u) > 0 \text{ and finite,}$$

$$\int_0^T\left|\left|\frac{2\alpha}{w^3 N}\sum_{i=1}^{N}\left[\xi_i^{\mu(s)}\left(\overline{\xi_i\Theta}+\sum_{j=1}^{N}\overline{\xi_i\xi_j}\tanh(y_j(\Gamma s))\right)\right]\right|\right|^2 d(\Gamma u) < \infty,$$

and

$$\int_0^T\left|\left|\frac{2\alpha^2}{w^6\Gamma N^2}\sum_{j=1}^{N}\left(\sum_{i=1}^{N}\xi_i^{\mu(t)}A_{ij}\right)^2\right|\right|^2 d(\Gamma u) \sim 0,$$

we have

$$0 < \int_0^T ||\gamma(\Gamma u)||^2\, d(\Gamma u) < \infty.$$

It follows that $\gamma(\Gamma t)$ satisfies the Novikov condition (see [17])

$$\mathbb{E}\left[\frac{1}{2}\int_0^T ||\gamma(\Gamma u)||^2 du\right] < \infty.$$

As a consequence we can apply Girsanov's Theorem (see [13], p.198); there exists a probability measure $\widetilde{\mathbb{P}}$ (risk neutral measure) under which $\widetilde{\mathbf{W}}(\Gamma t) = \int_0^t \gamma(\Gamma u)du - \mathbf{W}(\Gamma t)$ is an N-dimensional Brownian motion. It follows that

$$d(D(t)p(t)) = D(t)p(t)\frac{2\alpha}{w^3 N\Gamma}\sum_{i=1}^{N}\left(\xi_i^{\mu(t)}A_i\right)d\widetilde{\mathbf{W}}(\Gamma t),$$

and hence the discounted stock price is a $\widetilde{\mathbb{P}}$-martingale.

$\square$

By making the replacement $dW(\Gamma t) = \gamma(\Gamma t)dt - d\widetilde{W}(\Gamma t)$, dynamics (9) of the undiscounted stock price $p(t)$ becomes

$$\begin{aligned} p(t) &= p(0)\exp\left\{ \int_0^t R(s) - \frac{2\alpha^2}{w^6\Gamma N^2}\sum_{j=1}^N \left(\sum_{i=1}^N \xi_i^{\mu(t)} A_{ij}\right)^2 ds + \right. \\ & \left. \int_0^t \frac{2\alpha}{w^3 N\Gamma}\sum_{i=1}^N \left(\xi_i^{\mu(s)} A_i\right) d\widetilde{\mathbf{W}}(\Gamma s) \right\}. \end{aligned} \tag{10}$$

4.2 European call option pricing

Since, for $T - t = O(N)$, there exists a risk neutral measure $\widetilde{\mathbb{P}}$ for the discounted process $D(t)p(t)$, we can apply the usual scheme, relying on the martingale representation theorem, to show the existence of a replication portfolio for a derivative security pricing.

Consider $T - t = O(N)$; let $V(T)$ be an $\mathcal{G}_T$-measurable random variable representing the payoff at time T of a derivative security. The process $E(t) = \mathbb{E}_{\widetilde{\mathbb{P}}}[D(T)V(T)|\mathcal{G}_t]$ is a $\widetilde{\mathbb{P}}$-martingale (it follows from iterated conditioning); by the Martingale Representation Theorem there exists an initial capital $X(0)$ and a portfolio strategy $\Delta(t)$ such that $X(T) = V(T)$ almost surely and an adapted process $\phi(t)$ which constructs $E(t)$ out of $D(t)p(t)$. It follows that $D(t)X(t)$ is a $\widetilde{\mathbb{P}}$-martingale

$$D(t)X(t) = \mathbb{E}_{\widetilde{\mathbb{P}}}[D(T)X(T)|\mathcal{G}_t] = \mathbb{E}_{\widetilde{\mathbb{P}}}[D(T)V(T)|\mathcal{G}_t]. \tag{11}$$

The value $X(t)$ of the hedging portfolio in (11) is the capital needed at time t in order to successfully hedge the position in the derivative security with payoff $V(T)$. Hence $V(t)$ is the price of the derivative security at time t and we obtain the usual risk neutral pricing formula (see [19], p.218-222)

$$D(t)V(t) = \mathbb{E}_{\widetilde{\mathbb{P}}}[D(T)V(T)|\mathcal{G}_t], \quad 0 \le t \le T;$$

by recalling the definition of $D(t)$,

$$V(t) = \mathbb{E}_{\widetilde{\mathbb{P}}}[e^{-\int_t^T R(s)ds}V(T)|\mathcal{G}_t], \quad 0 \le t \le T.$$

Proposition 4.2 *Let us assume constant interest rate r; the price of a European call option with underlying $p(t)$ and strike K is*

$$c(t, p(t); K, r, \nu) = p(t)\psi(d(\theta, p(t))) - e^{-r\theta}K\psi\left(d\left(\theta, p(t)\right) - \sqrt{\nu}\right),$$

where $\theta = T - t$, $\nu = \frac{\alpha\sigma_N^2(T-t)}{w^2N^2}$, $d(\theta, p(t)) = \frac{\left[\log\frac{p(t)}{K} + \left(r + \frac{1}{2}\frac{\nu}{\theta}\right)\theta\right]}{\sqrt{\nu}}$ and ψ the erf function.

Proof

Consider the random variable

$$Y = \int_t^T \frac{2\alpha}{w^3N\Gamma}\sum_{i=1}^N \left(\xi_i^{\mu(s)}A_i\right) d\widetilde{\mathbf{W}}(\Gamma s) = \sum_{j=1}^N \int_t^T \frac{2\alpha}{w^3N\Gamma}\sum_{i=1}^N \left(\xi_i^{\mu(s)}A_{ij}\right) dW_j(\Gamma s).$$

Y has zero mean and, in the limit as N grows to infinity, finite variance ν (see prop. 3.1); since we consider N sufficiently large, we can assume $Var[Y] = \nu$. For N sufficiently large we can apply the results obtained at the end of proof of proposition (3.1); it follows that Y has a normal distribution with 0 mean and variance $\nu = \frac{\alpha\sigma_N^2(T-t)}{w^2N^2}$ and equation (10) becomes

$$p(T) = p(t)\exp\left\{\left(r - \frac{1}{2}\frac{\nu}{\theta}\right)\theta - \sqrt{\frac{\nu}{\theta}}\sqrt{\theta}Z\right\},$$

$\theta = T - t$ and $Z = -\frac{Y}{\sqrt{\nu}}$ is a standard normal random variable.

We can apply Black&Scholes formula (see [19], pp.218-220) and obtain the price of a call option with underlying p and strike K:

$$c(t, p(t); K, r, \nu) = p(t)\psi(d(\theta, p(t))) - e^{-r\theta}K\psi\left(d\left(\theta, p(t)\right) - \sqrt{\nu}\right),$$

where $d(\theta, p(t)) = \frac{\left[\log\frac{p(t)}{K} + \left(r + \frac{1}{2}\frac{\nu}{\theta}\right)\theta\right]}{\sqrt{\nu}}$ and ψ the *erf* function.

☐

5 Calibrating the MG

5.1 Calibration on the market

Minority Game is general purpose model, therefore, before calibrating it to price options, we first need to calibrate it to reflect the market properties. We review some basic fact about MG to better explain the calibration process. One of the most important variable of MG (for a detailed discussion see [9]) is σ^2.

Low value of σ^2 correspond to high market efficiency and vice-versa. As α (which depends on N and M, since $P = \frac{2^M}{N}$) decrease σ^2 decrease too, till it reach a minimum (see figure 1). We indicate with σ_c^2 this minimum and with α_c the corresponding phase transition point. Since σ_c^2 represent the maximum efficiency attainable in the market and efficiency is related with the information available, σ_c^2 can be thought as the point with the maximum information.

On the other side, volatility in option market is related with risk, which is linked with information too. The more the information available on the market the less the risk of an option and hence, the less the volatility.

The first calibration consists in linking this two quantity using the B&S equation derived in the previous section. Taking into account the expression of volatility ν and defining $\bar{\nu}^{mkt}$ as the minimum market volatility, we are looking for a value $\bar{w}$ such that

$$\sqrt{\bar{\nu}^{mkt}} = \sqrt{\alpha_c}\frac{\sigma_c}{\bar{w}N}. \tag{12}$$

The advantage of introducing the bet weight w, as is evident from equation (12), is to allow a meaningful calibration; without parameter w the values of σ_N required to match the options implied volatility would be extremely high implying either the choice of $\alpha < \alpha_c$, symmetric market phase known to be inefficient and not coherent with the real market, or $\alpha \gg \alpha_c$, game where agents play randomly.

The last step is to calibrate the MG on the options, i.e. we need to find the couple (α, σ_N^2) such that:

$$\sqrt{\nu^{impl}} = \sqrt{\alpha}\frac{\sigma_N}{\bar{w}N}$$

where $\sqrt{\nu^{impl}}$ is the option implied volatility available on market.



Figure 1: σ_N^2 as a function of α obtained numerically for different history lengths

5.2 Calibration on the options market

In the previous section we described the general procedure to calibrate the MG, here we implement it on DAX Index options. We choose the DAX Index since it is a total return index reinvesting dividends. The calibration of $\bar{w}$ is performed over VDAX Index, which is the index of implied volatility for DAX options. The minimum volatility, i.e. the minimum value of VDAX, since 03/01/2000 is reached the 11/02/2005 with a value of 10.98%. The next calibration, for each M, is performed for all options once. It is important to stress that it is not possible to calibrate the MG for M too small. The calibrated option value should correspond to a value α bigger than α_c which is around 0.34. Since the grid of values for α, given M, is fixed by the value of N which is an integer, it is possible to obtain only a fixed number of value for $\alpha > \alpha_c$. Consider $M = 3$, then $P = 8$ and for $N = 2$ we have that $\alpha = 4$. When $N = 21$ the value of α is 0.38. Therefore we have to find a minimum and calibrate it to match the value of options in 20 points. It is straightforward that we can not have an accurate calibration with this few points. The first M that allow a decent calibration is $M = 5$.

We choose 18 call options, with different maturities and different moneyness, and calibrate the MG over this option with M varying from 5 to 9 (in the tables only results for $M = 5$ and $M = 9$ are illustrated).In the head of the columns is indicated the maturity (expressed in years) and on the head of the rows is indicated the moneyness. For each maturity and moneyness, in the first row is indicated the market implied volatility $\sqrt{\nu^{impl}}$, in the second row are indicated the B&S volatility ν, the corresponding σ_N^2 and the α obtained from the calibration. Tables caption contains the corresponding M and the $\bar{w}$ obtained.

Comparison of the results shows that a better calibration is obtained as M increase. The values of the σ_N^2 are smaller and consequently the value of $\bar{w}$ are smaller for higher value of M. This is due to the fact that the more history M is used to construct the strategies, the better the strategies are and the market increases its efficiency and the

Moneyness	Maturity	0.18	0.52	1.01
1.0577	$\sqrt{\nu^{impl}}$	20.88%	21.02%	22.02%
	$\sigma^2_N \mid \sqrt{\nu}$	22.44% | 20.54%	22.44%| 20.54%	22.51% | 21.90%
	α	0.653	0.653	0.681
0.9967	$\sqrt{\nu^{impl}}$	17.66%	19.18%	20.72%
	$\sigma^2_N \mid \sqrt{\nu}$	20.65% | 17.52%	21.65% | 19.01%	22.44% | 20.54%
	α	0.604	0.627	0.653
0.9447	$\sqrt{\nu^{impl}}$	15.77%	18.10%	19.17%
	$(\sigma_N)^2 \mid \sqrt{\nu}$	19.32% | 16.03%	20.65% | 17.52%	21.65% | 19.01%
	α	0.582	0.604	0.627

Table 1: Calibration of MG with $M = 5$ and $\bar{w} = 0.03802281$.

Moneyness	Maturity	2.05	3.57	4.58
1.1062	$\sqrt{\nu^{impl}}$	23.79%	24.65%	25.58%
	$\sigma^2_N \mid \sqrt{\nu}$	23.93% | 24.11%	23.93% | 24.11%	25.14% | 26.45%
	α	0.711	0.711	0.653
0.9833	ν^{impl}	21.57%	22.79%	23.65%
	$\sigma^2_N \mid \sqrt{\nu}$	22.51% | 21.90%	22.51% | 21.90%	23.93% | 24.11%
	α	0.681	0.681	0.604
0.8849	$\sqrt{\nu^{impl}}$	19.61%	21.41%	22.41%
	$\sigma^2_N \mid \sqrt{\nu}$	21.65% | 19.01%	22.51% | 21.90%	22.51% | 21.90%
	α	0.627	0.681	0.582

Table 2: Calibration of MG with $M = 5$ and $\bar{w} = 0.03802281$.

value of σ^2_N decreases. On the other side the volatility of the real market is not changed and as a consequence we have a decrease of the weight $\bar{w}$ that each player gives to decisions.

6 Conclusions

In this paper we have shown that under the assumption of the MG with "fake" market histories as traders interaction model, the stock price dynamics follows the generalized exponential Brownian motion. This allowed us to apply the usual Black&Scholes pricing formula to value call European options and hence to calibrate, in terms of the game parameters α and w, the MG on the real options market.

The choice of a Minority Game with strategies $a^{\mu}_{s,i}$ uniform random variables taking values $\pm w$ reveals to be fundamental: the calibration of w on the minimum market volatility (the minimum value reached by VDAX Index in past years) allows to find values of the key game parameter α, calibrated on call options, greater than α_c (asymmetric game phase) and close to it (below 1 even with short history length M), corresponding to a game phase where agents coordinate in order to reach a state in which less resources are globally wasted. Using $w = 1$, that is the usual MG with $a^{\mu}_{s,i} = \pm 1$, and calibrating only on options, we get values of α far way from α_c and hence a game that is inefficient ($\alpha < \alpha_c$) or where agents play randomly ($\alpha \gg \alpha_c$).

The calibrated MG displays some interesting characteristics: as we already said it is

Moneyness	Maturity	0.18	0.52	1.01
1.0577	$\sqrt{\nu^{impl}}$	20.88%	21.02%	22.02%
	σ_N^2| $\sqrt{\nu}$	17.35% | 20.87%	17.48% | 21.02%	17.88% | 22.05%
	α	0.514	0.515	0.527
0.9967	$\sqrt{\nu^{impl}}$	17.66%	19.18%	20.72%
	σ_N^2| $\sqrt{\nu}$	15.97% | 17.74%	16.64% | 19.16%	17.42% | 20.85%
	α	0.474	0.492	0.513
0.9447	$\sqrt{\nu^{impl}}$	15.77%	18.10%	19.17%
	σ_N^2| $\sqrt{\nu}$	14.90% | 15.76%	16.07% | 18.09%	16.64% | 19.16%
	α	0.448	0.479	0.492

Table 3: Calibration of MG with $M = 9$ and $\bar{w} = 0.00143437$.

Moneyness	Maturity	2.05	3.57	4.58
1.1062	$\sqrt{\nu^{impl}}$	23.79%	24.65%	25.58%
	σ_N^2| $\sqrt{\nu}$	18.66% | 23.84%	18.97% | 24.66%	19.28% | 25.53%
	α	0.548	0.557	0.567
0.9833	$\sqrt{\nu^{impl}}$	21.57%	22.79%	23.65%
	σ_N^2| $\sqrt{\nu}$	17.66% | 21.57%	18.18% | 22.79%	18.53 | 23.68%
	α	0.521	0.536	0.546
0.8849	$\sqrt{\nu^{impl}}$	19.61%	21.41%	22.41%
	σ_N^2| $\sqrt{\nu}$	16.78% | 19.63%	17.62% | 21.42%	17.96% | 22.41%
	α	0.499	0.520	0.532

Table 4: Calibration of MG with $M = 9$ and $\bar{w} = 0.00143437$.

asymmetric and with α close to α_c; moreover, as it is evident from the tables displaying calibration results, a better calibration is obtained as M increases. The values of the σ_N^2 are smaller and consequently the value of $\bar{w}$ are smaller for higher values of M; when the market efficiency improves each agent decreases the weight of its buy or sell decision: players acting in not efficient markets characterized by scarce liquidity are required to increase the weight of the bets, like it happens in real market.

References

[1] Bouchaud J.P. and Potters M. (2000) *Theory of financial risks: from statistical physics to risk management.* Cambridge University Press, Cambridge.

[2] Cavagna A., Garrahan J P., Giardina I. and Sherrington D. (1999) A thermal model for adaptive competition in a market. *Phys. Rev. Lett.*, **83**, 4429.

[3] Challet D., Chessa A., Marsili M. and Zhang Y. C. (2001) From Minority Games to real markets. *Quantitative Finance*, **1**, 168–176.

[4] Challet D. and Zhang Y. C. (1997) Emergence of cooperation and organization in an evolutionary game. *Physica A*, **246**, 407.

[5] Challet D., Marsili M. and Zhang Y. C. (2001) Stylized facts of financial markets and market crashes in Minoriy Games *Physica A*, **294**, 514–524.

[6] Coolen A.C.C. (2005) *The Mathematical Theory of Minority Games - Statistical Mechanics of Interacting Agents.* Oxford University Press.

[7] Coolen A.C.C. and J.A.F. Heimel (2001) Dynamical solution of the on-line minority game. *J. Phys. A*, **34**, 10783–10804.

[8] Coolen A.C.C. (2005) Generating functional analysis of minority games with real market histories. *J. Phys. A*, **38**, 2311–2347.

[9] CHALLET D., MARSILI M. AND ZHANG Y. C. (2004) *Minority Games Interacting agents in financial markets.* Oxford Finance Series.

[10] Giardina I., Bouchaud J.P. and Mezard M. (2001) Microscopic models for long ranged volatility correlations. *Physica A*, **299**, 28-39.

[11] Jefferies P., Hart M.L., Hui P.M. and Johnson N.F. (2001) From market games to real-world markets. *Eur.Phys. J. B* **20**, 493-501.

[12] Johnson N.F., Hart M., Hui P.M. and Zheng D (2000) Trader dynamics in a model market. *J. Theo. App. Fin.*, **3**, 443.

[13] Karatzas I. and Shreve S.E. (1991) *Brownian Motion and Stochastic Calculus.* Springer-Verlag, New York.

[14] Lux T. and Marchesi M. (1999) Scaling and criticality in a stochastic multi-agent model of financial market. *Nature*, **397**, 498-500.

[15] Marsili M. and Challet D. (2001) Continuum time limit and stationary states of the minority game. *Phys. Rev. E*, **6405 (5)**, art. no.056138, Part 2.

[16] Moro E. (2004) Minority Game: an introductory guide. *Advances in Condensed Matter and Statistical Physics.* Nova Science Publishers, Madrid.

[17] Novikov A.A. (1971) On moment inequalities for stochastic integrals. *Theory Prob. Appl.*, **17**, 717-720.

[18] Ortisi M. (2008) Polynomial-rate convergence to the stationary state for the continuum-time limit of the Minority Game. *Journal of Applied Probability*, **45**, 376-387.

[19] Shreve S.E. (2004) *Stochastic Calculus for Finance, V. II.* Springer Finance, NY.

[20] Stanley H.S. and Mantegna R.N. (2000) *An introduction to econophysics: correlations and complexity in finance.* Cambridge University Press, Cambridge.